

PRIMITIVE ROOTS IN QUADRATIC FIELDS II

JOSEPH COHEN

ABSTRACT. This paper is continuation of the paper "Primitive roots in quadratic field". We consider an analogue of Artin's primitive root conjecture for algebraic numbers which is not a unit in real quadratic fields. Given such an algebraic number, for a rational prime p which is inert in the field the maximal order of the unit modulo p is $p^2 - 1$. An extension of Artin's conjecture is that there are infinitely many such inert primes for which this order is maximal. we show that for any choice of 85 algebraic numbers satisfying a certain simple restriction, there is at least one of the algebraic numbers which satisfies the above version of Artin's conjecture.

1. INTRODUCTION

This paper is continuation of the paper "Primitive roots in quadratic field" ([8]). In this paper we considered an analogue of Artin's primitive root conjecture for units in real quadratic fields. Given such a nontrivial unit, for any rational prime p which is inert, the maximal order of the unit modulo p is $p + 1$. An extension of Artin's conjecture is that there are infinitely many such inert primes for which this order is maximal. This is known at present only under the Generalized Riemann Hypothesis. Unconditionally, we showed that for any choice of 7 units in different real quadratic fields satisfying a certain simple restriction, there is at least one of the units which satisfies the above version of Artin's conjecture. In this paper we want to extend this result for any algebraic integer modulo inert prime p . we will prove

Theorem 1.1. *Let $K = \mathbb{Q}(\sqrt{\Delta})$ be a quadratic field and $\{\alpha_i\}_{i=1}^{85}$ be a set of 85 integers of K such that*

- (1) *The norms $N(\alpha_i) = \alpha_i \sigma(\alpha_i)$, of the α_i 's, are multiplicatively independent.*

Date: February 1, 2008.

In partial fulfillment for the Ph.D. degree. Supported by grants from the Technion-Israel Institute of Technology.

(2) $5N(\alpha_i)\Delta$, $N(\alpha_i)$ are not perfect squares.

(3) $M(\alpha_i) = \sigma(\alpha_i)/\alpha_i$ are multiplicatively independent.

Then at least one of the 85 integers has at least order $\frac{p^2-1}{24} \pmod{p}$ for infinitely many inert primes p in K .

Note that in the case of split primes, Narkiewicz ([12]) proved a much stronger result.

Since in our case the order is $p^2 - 1$, which is not “linear”, some of their divisors are too big and we cannot use the method of [6]. But since $p^2 - 1 = (p - 1)(p + 1)$ can be factored into two linear factors, with the following remark we can still use the method of [6]

Remark 1.2. Consider an algebraic number α in $K = \mathbb{Q}(\sqrt{\Delta})$. Let $p \nmid \alpha$ be an inert prime in K . Since

$$(1) M(\alpha) \equiv \alpha^{p-1} \pmod{(p)},$$

$$(2) N(\alpha) \equiv \alpha^{p+1} \pmod{(p)},$$

we have:

$$\text{ord}(M(\alpha)) \mid \text{ord}(\alpha) \pmod{(p)} \text{ and } \text{ord}(N(\alpha)) \mid \text{ord}(\alpha) \pmod{(p)}$$

In addition:

$$\text{ord}(M(\alpha)) \mid p + 1 \text{ and } \text{ord}(N(\alpha)) \mid p - 1$$

But,

$$\left(\frac{p-1}{2}, p+1\right) = 1 \text{ or } \left(p-1, \frac{p+1}{2}\right) = 1.$$

So,

$$\text{ord}(M(\alpha)) \text{ord}(N(\alpha)) \mid 2 \text{ord}(\alpha) \pmod{(p)}$$

Let e_1 and e_2 be some integers. If we prove that $M(\alpha)$ and $N(\alpha)$ have simultaneously at least orders $\frac{p+1}{e_1}$ and $\frac{p-1}{e_2}$, respectively, then we will obtain that α has at least order $\frac{p^2-1}{2e_1e_2}$. This way we reduce the problem to a “linear” problem.

1.1. Notation and Preliminaries. Let $\pi(y; m, s)$ the number of primes $p \leq y$ such that $p \equiv s \pmod{m}$ where m and s are some integers, and

$$E(y; m, s) := \pi(y; m, s) - \frac{Li(y)}{\varphi(m)}$$

where $Li(y) = \int_2^y \frac{dt}{\log t}$. Also set

$$E(x; m) := \max_{1 \leq y \leq x} \max_{(s, m)=1} |E(y; m, s)|.$$

Denote $\mathcal{A} = \{p^2 - 1 | p \leq x, p \equiv u \pmod{v}\}$ where u, v are some given integers such that $(u, v) = 1$, and take $X = \frac{Li(x)}{\varphi(v)}$.

For a square-free integer d , $(d, v) = 1$, denote

$$\begin{aligned} |\mathcal{A}_d| &:= |\{a \in \mathcal{A} : a \equiv 0 \pmod{d}\}| \\ &= |\{p^2 - 1 : p \leq x, p \equiv u \pmod{v}, p^2 - 1 \equiv 0 \pmod{d}\}| \\ &= \sum_{\substack{m=1 \\ m^2-1 \equiv 0 \pmod{d}}}^d |\{p | p \leq x, p \equiv u \pmod{v}, p \equiv m \pmod{d}\}| \\ &= \sum_{\substack{m=1 \\ m^2-1 \equiv 0 \pmod{d} \\ (m, d)=1}}^d |\{p | p \leq x, p \equiv u \pmod{v}, p \equiv m \pmod{d}\}| \end{aligned}$$

By the Chinese Remainder Theorem, for each m there exists an integer l_m such that

$$|\mathcal{A}_d| = \sum_{\substack{m=1 \\ m^2-1 \equiv 0 \pmod{d} \\ (m, d)=1}}^d |\{p | p \leq x, p \equiv l_m \pmod{dv}\}|;$$

Since $|\{p | p \leq x, p \equiv l_m \pmod{dv}\}|$ is asymptotically independent of m , there exists some integer l such that

$$|\mathcal{A}_d| = \pi(x; dv, l) \sum_{\substack{m=1 \\ m^2-1 \equiv 0 \pmod{d} \\ (m, d)=1}}^d 1 = \pi(x; dv, l) \rho(d)$$

$$\text{where } \rho(d) = \sum_{\substack{m=1 \\ m^2-1 \equiv 0 \pmod{d} \\ (m,d)=1}}^d 1.$$

We note that $\rho(q) = 2$ for any prime q , and hence for any square-free d , $\rho(d) = 2^{\nu(d)}$ where $\nu(d)$ denotes the number of prime divisors of d .

By the definition of $E(x; dv, l)$,

$$|\mathcal{A}_d| = \frac{\rho(d)}{\varphi(d)} \frac{Lix}{\varphi(v)} + \rho(d) E(x; dv, l) = \frac{2^{\nu(d)}}{\varphi(d)} X + 2^{\nu(d)} E(x; dv, l).$$

For any prime q define $\omega(q) := \frac{2q}{\varphi(q)}$, $\omega(d) = \prod_{q|d} \omega(q) = \frac{2^{\nu(d)} d}{\varphi(d)}$ and

$$R_d := |\mathcal{A}_d| - \frac{\omega(d)}{d} X = 2^{\nu(d)} E(x; dv, l)$$

Finally, we define the Möbius function, $\mu(1) = 1$ and for a square-free $d = p_1 \cdots p_k$, $\mu(d) = (-1)^k$.

Now we want to prove two lemmas.

Lemma 1.3. *For any prime $q > 3$ which is relatively prime to v we have:*

$$(1.1) \quad 0 \leq \frac{2}{q-1} \leq \frac{1}{2}.$$

$$(1.2) \quad \sum_{w \leq q < z} \frac{2}{q-1} \log q - 2 \log \frac{z}{w} = O(1) \quad (2 \leq w \leq z)$$

where O does not depend on z or w .

$$(1.3) \quad \prod_{\substack{2 < q < z \\ q \nmid v}} \left(1 - \frac{2}{q-1}\right) \gg \frac{1}{\log^2 z}.$$

Proof. Since $q > 3$, it is clear that (1.1) holds.

As for the second equality, $\sum_{w \leq q < z} \frac{2}{q-1} \log q = 2 \sum_{w \leq q < z} \frac{\log q}{q} \frac{q}{q-1} = 2 \sum_{w \leq q < z} \frac{\log q}{q} (1 + \frac{1}{q-1}) = 2 \sum_{w \leq q < z} \frac{\log q}{q} + 2 \sum_{w \leq q < z} \frac{\log q}{q(q-1)} = 2 \log \frac{z}{w} + O(1) \quad (\sum_{p < x} \frac{\log p}{p} = \log x + O(1)).$

Hence we get (1.2). Finally,

$$\begin{aligned}
\prod_{\substack{2 < q < z \\ q \nmid v}} \left(1 - \frac{2}{q-1}\right) &\gg \prod_{2 < q < z} \left(1 - \frac{2}{q-1}\right) \\
&= \exp\left(\log \prod_{2 < q < z} \left(1 - \frac{2}{q-1}\right)\right) \\
&= \exp\left(\sum_{2 < q < z} \log\left(1 - \frac{2}{q-1}\right)\right) \\
&\gg \exp\left(\sum_{2 < q < z} \left(-\frac{2}{q-1} - \frac{4}{(q-1)^2}\right)\right)
\end{aligned}$$

Since

$$\frac{2}{q-1} = \frac{2}{q} + \frac{2}{q(q-1)} \leq \frac{2}{q} + \frac{2}{(q-1)^2}$$

and $\sum_{2 < q < z} \frac{2}{(q-1)^2}$ converges, we get

$$\prod_{2 < q < z} \left(1 - \frac{2}{q-1}\right) \gg \exp\left(-\sum_{2 < q < z} \frac{2}{q}\right)$$

Since

$$\sum_{2 < q < z} \frac{2}{q} \sim 2 \log \log z$$

we have

$$\exp\left(-\sum_{2 < q < z} \frac{2}{q}\right) \gg \exp(-2 \log \log z) = \frac{1}{\log^2 z}$$

□

Lemma 1.4. *For any square-free natural number d , $(d, v) = 1$, and a real number $A > 0$, there exist constants $c_2(\geq 1)$ and $c_3(\geq 1)$ such that*

$$(1.4) \quad \sum_{d < \frac{X^{\frac{1}{2}}}{(\log x)^{c_2}}} \mu^2(d) 3^{\nu(d)} |R_d| \leq c_3 \frac{X}{\log^A X}, \quad (X \geq 2)$$

Proof. Denote by S_R the term which we need to estimate.

$$S_R = \sum_{d < \frac{X^{\frac{1}{2}}}{(\log x)^{c_2}}} \mu^2(d) 3^{\nu(d)} |R_d|$$

By the definitions of R_d and $E(x; dv)$

$$S_R \leq \sum_{d < \frac{X^{\frac{1}{2}}}{(\log x)^{c_2}}} \mu^2(d) 6^{\nu(d)} |E(x; dv)|.$$

Since $E(x; dv) \ll \frac{x}{dv}$ if $d \leq \frac{x}{v}$, we get that

$$S_R \ll x^{\frac{1}{2}} \sum_{d < \frac{X^{\frac{1}{2}}}{(\log x)^{c_2}}} \frac{\mu^2(d) 6^{\nu(d)}}{d^{\frac{1}{2}}} |E(x; dv)|^{\frac{1}{2}}.$$

By Cauchy's Inequality,

$$S_R \ll x^{\frac{1}{2}} \left(\sum_{d < X^{\frac{1}{2}}} \frac{\mu^2(d) 6^{2\nu(d)}}{d} \right)^{\frac{1}{2}} \left(\sum_{dv < \frac{vX^{\frac{1}{2}}}{(\log x)^{c_2}}} |E(x; dv)| \right)^{\frac{1}{2}}.$$

For sufficiently large x we obtain

$$S_R \ll x^{\frac{1}{2}} \left(\sum_{d < x^{\frac{1}{2}}} \frac{\mu^2(d) 6^{2\nu(d)}}{d} \right)^{\frac{1}{2}} \left(\sum_{dv < \frac{x^{\frac{1}{2}}}{(\log x)^{c_2}}} |E(x; dv)| \right)^{\frac{1}{2}}.$$

With Bombieri-Vinogradov Theorem [2] (given any positive constant e_1 , there exists a positive constant e_2 such that $\sum_{d < \frac{x^{\frac{1}{2}}}{\log^{e_2} x}} E(x; d) =$

$O(\frac{x}{\log^{e_1} x})$) for the last sum and the inequality $\sum_{d < w} \frac{\mu^2(d) 36^{\nu(d)}}{d} \leq (\log w + 1)^{36}$, [5], p.115, equation (6.7)) we find that for given constant B there exists c_2 such that

$$S_R \ll \frac{x}{\log^B x}.$$

So, for given A there exists c_2 such that

$$S_R \ll \frac{X}{\log^A X}$$

where $\ll$ depends on v and c_2 . □

1.2. Proof of Theorem 1.1 – the sieve part. In this section we use the Selberg lower bound sieve and show that there is some small real number δ_1 and some constant $c(\delta_1) > 0$ (which depends on δ_1) such that for at least $c(\delta_1) \frac{x}{\log^3 x}$ primes $p \leq x$, $p \equiv u \pmod{v}$, if $q|p^2 - 1$ then either $q > x^{1/8+\delta_1}$ or $q|v$.

Now, define $S(\mathcal{A}, z, v) = |\{a | a \in \mathcal{A}, (a, \prod_{\substack{p < z \\ p \nmid v}} p) = 1\}|$ and define a function g by $g(t_0) = 1, t_0 = 4.42$ and $g(t) < 1$ for $t > t_0$. Then (see [5, Theorem 7.4, page 219]):

Lemma 1.5. *We have*

$$(1.5) \quad S(\mathcal{A}, z, v) \geq X \prod_{\substack{q < z \\ q \nmid v}} (1 - \frac{\omega(q)}{q}) \{1 - g(\frac{\log X}{2 \log z}) + O(\frac{(\log \log 3X)^8}{\log X})\}$$

where the O -term does not depend on X or on z .

By Lemmas 1.3 and 1.4, (1.1), (1.2) and (1.4) hold. Hence we can use Lemma 1.3 with $z = X^{\frac{1}{8} + \delta_0}$

$$S(\mathcal{A}, X^{\frac{1}{8} + \delta_0}, v) \geq X \prod_{\substack{q < X^{\frac{1}{8} + \delta_0} \\ q \nmid v}} (1 - \frac{2}{q-1}) \{1 - g(\frac{1}{2} \frac{\log X}{\log X^{\frac{1}{8} + \delta_0}}) + O(\frac{(\log \log 3X)^8}{\log X})\}.$$

By Lemma 1.3 (1.3) we have for δ_0 sufficiently small

$$S(\mathcal{A}, X^{\frac{1}{8} + \delta_0}, v) \gg \frac{X}{\log^2 X} \gg \frac{x}{\log^3 x}$$

Thus for such δ_0 we obtain that there is a constant $c(\delta_0) > 0$ (which depends on δ_0) such that for at least $c(\delta_0) \frac{x}{\log^3 x}$ primes $p \leq x$, $p \equiv u \pmod{v}$ if $q | p^2 - 1$ then either $q > x^{1/8 + \delta_0}$ or $q | v$. Hence we obtain that for all $0 < \delta_1 < \delta_0$ there is a constant $c(\delta_1) > 0$ (which depends on δ_1) such that for at least $c(\delta_1) \frac{x}{\log^3 x}$ primes $p \leq x$, $p \equiv u \pmod{v}$, if $q | p^2 - 1$ then either $q > x^{1/8 + \delta_1}$ or $q | v$.

1.3. Proof of Theorem 1.1 – The algebraic part.

1.3.1. *Construction of the arithmetic sequence.* Let $K = \mathbb{Q}(\sqrt{\Delta})$ be any quadratic field, $\mathcal{O}$ the integers ring of K , $\alpha \in \mathcal{O}$ any algebraic integer and $a = N(\alpha)$. In this section we want to construct integers u and v , $(u, v) = 1$ such that for all primes p such that $p \equiv u \pmod{v}$, the discriminant Δ of $\mathbb{Q}(\sqrt{\Delta})$ and a satisfy

$$\left(\frac{\Delta}{p}\right) = \left(\frac{a}{p}\right) = -1.$$

This means that p is inert and a is not a quadratic residue $(\text{mod } p)$. In addition we want to obtain by the construction that $(\frac{p^2-1}{24}, v) = 1$ (since after sieving the small factors of $\frac{p^2-1}{24}$ we may be left with small factors which divide v , see previous section).

In order to fulfill these demands, we will first show that there exist infinitely many primes p satisfying the following simultaneous conditions

$$(1.6) \quad \left(\frac{-1}{p}\right) = \left(\frac{5}{p}\right) = \left(\frac{a}{p}\right) = \left(\frac{\Delta}{p}\right) = -1.$$

This condition is equivalent to the condition:

$$B(p) = \left(1 - \left(\frac{-1}{p}\right)\right)\left(1 - \left(\frac{5}{p}\right)\right)\left(1 - \left(\frac{a}{p}\right)\right)\left(1 - \left(\frac{\Delta}{p}\right)\right) \neq 0$$

Since the Legendre symbol is a multiplicative function, we obtain,

$$\left(1 - \left(\frac{-1}{p}\right)\right)\left(1 - \left(\frac{a}{p}\right) - \left(\frac{\Delta}{p}\right) + \left(\frac{a\Delta}{p}\right) - \left(\frac{5}{p}\right) + \left(\frac{5a}{p}\right) + \left(\frac{5\Delta}{p}\right) - \left(\frac{5a\Delta}{p}\right)\right).$$

Let S be the set of all integers of the form $n = (-1)^{b_0} 5^{b_1} a^{b_2} \Delta^{b_3}$, $b_i \in \{0, 1\}$. Then

$$(1.7) \quad \sum_{p \leq Z} B(p) = \sum_{n \in S} (-1)^{b_0+b_1+b_2+b_3} \sum_{p \leq Z} \left(\frac{n}{p}\right), \quad b_i \in \{0, 1\}.$$

By the assumption in the theorem each $n \in S$ is not a perfect square when $\sum_{i=0}^3 b_i$ is odd.

This assumption, together with the fact that for n not a perfect square (by the reciprocity law for Legendre symbol)

$$\sum_{p \leq Z} \left(\frac{n}{p}\right) = o(\pi(Z)) \quad \text{as } Z \rightarrow \infty$$

implies that $\sum_{p \leq Z} B(p)$ is asymptotic to at least $\pi(Z)$ (since all the negative summands contribute $o(\pi(Z))$ and at least the natural number 1 contributes $\pi(Z)$). This shows that the simultaneous conditions have infinitely many solutions p .

We fix some particular p_0 satisfying the condition (1.6) and for each odd prime $l \neq 3$, such that $l \nmid 24a\Delta$, we define $u_l = p_0$ if $l \nmid p_0^2 - 1$, and $u_l = 9p_0$ otherwise.

Claim 1.6. $l \nmid u_l^2 - 1$.

Proof. If $u_l = p_0$ then by the assumption $l \nmid p_0^2 - 1$, so $l \nmid u_l^2 - 1$. If $u_l = 9p_0$, assume, by reductio ad absurdum, that $l \mid u_l^2 - 1$. Hence $l \mid 81p_0^2 - 1$. Since, $l \mid p_0^2 - 1$, we obtain that $l \mid 80p_0^2$. On the other hand, by our condition, $\left(\frac{5}{p_0}\right) = -1$ so $\left(\frac{p_0}{5}\right) = -1$ ($p_0 \equiv 1 \pmod{4}$). Hence $p_0 \equiv 2$ or $3 \pmod{5}$. Since $l \mid p_0^2 - 1$ and $p_0 \equiv 2$ or $3 \pmod{5}$ we conclude that $l \nmid 5$. Using the assumption that $l \mid p_0^2 - 1$ we deduce that $l \neq p_0$ (if $l = p_0$ then $l \nmid p_0^2 - 1$). Hence $(l \neq 2, 3) \mid 80p_0^2$, which is a contradiction. $\square$

In addition let $u_2 = p_0$ if $8 \mid p_0^2 - 1$ and $u_2 = p_0 - 8$ if $16 \mid p_0^2 - 1$. Likewise we take $u_3 = p_0$ if $3 \mid p_0^2 - 1$ and $u_3 = p_0 - 3$ if $9 \mid p_0^2 - 1$.

Let $v = 24a\Delta$ and u be the common solution of $u \equiv u_2 \pmod{16}$, $u \equiv u_3 \pmod{9}$ and all the congruences $u \equiv u_l \pmod{l}$. Such a solution exists, by the Chinese Remainder Theorem.

Since $l \nmid u^2 - 1$ for every odd prime $l \neq 2, 3$, $l \mid v$, and by the construction $(\frac{u^2-1}{24}, 6) = 1$ we conclude that $(\frac{u^2-1}{24}, v) = 1$.

Finally, if $p \equiv u \pmod{v}$, then $p \equiv p_0 \pmod{24}$ and $p \equiv p_0$ or $4p_0 \pmod{l}$ for all odd primes $l \mid v$. So, $(\frac{\Delta}{p}) = (\frac{\Delta}{p_0}) = -1$, and similarly for a . This completes the construction of u and v .

Note that by the construction of the integers u and v we have that $(u, v) = 1$. (take l an odd prime number, $l \mid v = 24a\Delta$ and assume that $l \mid u$. Since $u \equiv u_l \pmod{l}$, $l \mid u_l$. Hence $l \mid p_0$ or $9p_0$ (in this case $l \neq 2, 3$). In other words $l = p_0$. But $p_0 \nmid 24a\Delta$ (p_0 fulfills the simultaneous condition (1.6)) and $l \mid 24a\Delta$).

1.3.2. The last step of the proof. As we saw at the previous subsections, for at least $c(\delta_1) \frac{x}{\log^3 x}$ primes $p \leq x$, $p \equiv u \pmod{v}$, if $q \mid p^2 - 1$ then $q > x^{1/8+\delta_1}$ or $q \mid v$. Since $(\frac{p^2-1}{24}, v) = 1$, if $q \mid \frac{p^2-1}{24}$ then $q > x^{1/8+\delta_1}$.

Since by the construction of u and v , $p \equiv 1 \pmod{4}$ and $(\frac{p^2-1}{24}, v) = 1$, we have that $\frac{p-1}{4}$ and $\frac{p+1}{2}$ are odd. In addition if $p \equiv 1 \pmod{3}$ then $(\frac{p-1}{12}, v) = 1$ and if $p \equiv -1 \pmod{3}$ then $(\frac{p+1}{6}, v) = 1$.

If we conclude the result about $p - 1$ and $p + 1$ we have for at least $c_1(\delta_1) \frac{x}{\log^3 x}$ primes $p \leq x$, $p \equiv u \pmod{v}$, if $q \mid \frac{p-1}{d_-}$ or $q \mid \frac{p+1}{d_+}$ then $q > x^{1/8+\delta_1}$, where $d_- = 4$ or 12 and $d_+ = 6$ or 2 , respectively.

For the last step of the proof we need to use a version of Lemma 4 from Narkiewicz [12], which generalized Lemma 2 in [3].

Lemma 1.7. *If $a_1, \dots, a_k$ are multiplicatively independent algebraic numbers of an algebraic number-field K , G the subgroup of $K^\star$ generated by $a_1, \dots, a_k$, and for any prime ideal $\mathbf{P}$ not dividing $a_1 \cdots a_k$ we denote by $G_{\mathbf{P}}$ the reduction of $G \pmod{\mathbf{P}}$, then for all positive y one can have $|G_{\mathbf{P}}| < y$ for at most $O(y^{1+\frac{1}{k}})$ prime ideals $\mathbf{P}$, with the implied constant being dependent on the a_i 's and K .*

Proof. According to [12],

For any real number T denote by $M = M(T)$ the set of all k -element sequences $(r_1, \dots, r_k)$ of non-negative integers satisfying

$$|r_1| + |r_2| + \dots + |r_k| \leq T$$

It is easy to see that for T tending to infinity $|M(T)| = (c + o(1))T^k$ with suitable positive constant $c = c_k$. If now P is a prime ideal for which $|G_P| < y$, then select T to be the smallest rational integer with $cT^k > 2y$ and let $a_i = \frac{b_i}{c_i}$ for $i = 1, \dots, k$. There exist two distinct sequences $Z = (z_i)$, $W = (w_i)$ in $M(T)$ for which

$$P \mid \frac{b_1^{z_1} \dots b_k^{z_k}}{c_1^{z_1} \dots c_k^{z_k}} - \frac{b_1^{w_1} \dots b_k^{w_k}}{b_1^{w_1} \dots c_k^{w_k}}$$

Hence for sufficient large P (since the a_i 's are multiplicatively independent).

$$P \mid \frac{b_1^{z_1} \dots b_k^{z_k} - b_1^{w_1} \dots b_k^{w_k}}{c_1^{[z_1, w_1]} \dots c_k^{[z_k, w_k]}} = D \neq 0$$

Thus for sufficient large P

$$\nu_P(\Pi a_i^{z_i - w_i} - 1) \geq 0$$

where ν_P denotes the P -adic valuation and it follows that for fixed $z_1 - w_1, \dots, z_k - w_k$ we obtain $\ll \log(\max_j |\bar{a}_j|^{2T}) \ll T$ possibilities for P . Finally we obtain

$$|\{P \mid |G_P| < y\}| \ll T^{1+k} \ll y^{1+\frac{1}{k}}$$

□

Look at the $p - 1$ case (the case of $p + 1$ is similar). We have for at least $c_1(\delta_1) \frac{x}{\log^3 x}$ primes $p \leq x$, $p \equiv u \pmod{v}$ such that if $q \mid \frac{p-1}{d_-}$ then $q > x^{1/8+\delta_1}$ where $d_- = 4$ or 12 . Let $p - 1 = d_- q_1(p) q_2(p) \dots q_m(p)$, $q_m(p) > q_{m-1}(p) > \dots > q_1(p)$, $m \leq 7$, and let a be some integer where $\bar{a}$ its image in $\mathbb{F}_p^*$.

Denote by S_n the set $S_n = \{a_1, \dots, a_n\}$ where $a_1, \dots, a_n$ are multiplicatively independent integers and take seven integers $a_{i_1}, \dots, a_{i_7}$ from S_n and assume that at least one prime, say $q_1(p)$, which is greater than $x^{1/8+\delta_1}$, divides $[\mathbb{F}_p^* : \langle \bar{a}_{i_k} \rangle]$ for $k = 1, \dots, 7$.

Then $|\langle \bar{a}_{i_k} \rangle| = \frac{p-1}{[\mathbb{F}_p^* : \langle \bar{a}_{i_k} \rangle]} \ll p^{7/8-\delta_1} \ll x^{7/8-\delta_1}$, $k = 1, \dots, 7$. Since $q_1(p)$ divides $[\mathbb{F}_p^* : \langle \bar{a}_{i_k} \rangle]$ for $k = 1, \dots, 7$ and $\mathbb{F}_p^*$ is a cyclic group, $|\langle a_{i_1}, \dots, a_{i_7} \rangle| \bmod p \ll x^{7/8-\delta_1}$. By Lemma 1.7 it occurs in at most $O((x^{7/8-\delta_1})^{8/7}) = O(x^{1-8/7\delta_1})$ primes $p \leq x$ which is a negligible number relatively to $c_1(\delta_1) \frac{x}{\log^3 x}$ for sufficiently small δ_1 .

So, for at most six integers from $a_{i_1}, \dots, a_{i_7}$, $q_1(p)$ divides $[\mathbb{F}_p^* : \langle \bar{a}_{i_k} \rangle]$ for $k = 1, \dots, 7$. In other words for at most six from $a_{i_1}, \dots, a_{i_7}$ $q_1(p)$ does not divide $|\langle \bar{a}_{i_k} \rangle|$. Hence for at least one integer, say a_n , $q_1(p)$ divide $|\langle \bar{a}_n \rangle|$.

Denote by S_{n-1} the set $\{a_1, \dots, a_{n-1}\}$. By repeating the former process for S_{n-1} we obtain that for at least one integer, say a_{n-1} , $q_1(p)$ divide $|\langle \bar{a}_{n-1} \rangle|$.

We continue this process till we obtain the set $T_1 = \{a_7, \dots, a_n\}$ where $a_7, \dots, a_n$ are multiplicatively independent integers such that $q_1(p)$ divides $|\langle \bar{a}_t \rangle|$ for $t = 7, \dots, n$.

By repeating this process for $q_2(p)$ we obtain for the set $T_2 = \{a_{13}, \dots, a_n\}$ that $q_2(p)$ divides $|\langle \bar{a}_t \rangle|$ for $t = 13, \dots, n$.

Again, by repeating this process for $q_m(p)$ we obtain for the set $T_m = \{a_{6m+1}, \dots, a_n\}$ that $q_m(p)$ divides $|\langle \bar{a}_t \rangle|$ for $t = 6m+1, \dots, n$.

Since the maximum value of m is 7, if we take $n = 6m+1 = 43$ multiplicatively independent integers we obtain that one of them have at least the order $\frac{p-1}{d_-}$.

Let us look on the algebraic number $M(\alpha)$. By the same method we obtain that one of 43 $M(\alpha)$ have at least the order $\frac{p+1}{d_+}$. one of $42 + 43 = 85$ has, at least, the order $\frac{p^2-1}{24}$. This completes the proof of Theorem 1.1

Acknowledgement I wish to gratefully acknowledge my Ph.D supervisors Prof. Zee'v Rudnick and Prof. Jack Sonn for the helpful suggestion and the fruitful ideas, especially for Remark 1.2.

REFERENCES

- [1] E. Artin, *Collected Papers*, Reading, MA: Addison-Wesley (1965).

- [2] E. Bombieri, *On the large sieve*, Mathematika 12 (1965), 201-225.
- [3] R. Gupta and R. Murty, *A remark on Artin's conjecture*, Invent. Math. 78, 127-130 (1984).
- [4] R. Gupta, V. Kumar Murty and M. Ram Murty, *The Euclidean algorithm for S integers*, CNS Conference Proceedings, Vol.7 (1985), 189-202.
- [5] Halberstam and Richert, *Sieve Methods*, Academic Press, London 1974.
- [6] D. R. Heath-Brown, *Artin's conjecture for primitive roots*, Quart. J. Math. Oxford (2), 37 (1986), 27-38.
- [7] C. Hooley, *On Artin's Conjecture*, J. Reine Angew. Math. 226 (1967), 209-220.
- [8] J. Cohen, *Primitive Roots in Quadratic Fields*,
<http://arxiv.org/abs/math.NT/0310205>.
- [9] P. Kurlberg and Z. Rudnick, *Hecke theory and equidistribution for the quantization of linear maps of the torus*, Duke Math. Jour. 103 (2000), 47-77.
- [10] P. Kurlberg and Z. Rudnick, *On quantum ergodicity for linear maps of the torus*, Comm. in Math. Physics. 222 (2001) 1, 201-227.
- [11] D. Marcus, *Number Fields*, Springer, New York 1977.
- [12] W. Narkiewicz, *A note on Artin's conjecture in algebraic number fields*, J. Reine Angew. Math. 381 (1987), 110-115.
- [13] W. Narkiewicz, *Units in residue classes*, Arc. Math., Vol. 51, 238-241 (1988)
- [14] O. Taussky, *Introduction into connections between algebraic number theory and integral matrices*, Appendix to H. Cohn *A Classical Invitation to Algebraic Numbers and Class Fields*, Springer, New York 1978.

E-mail address: `coheny@techunix.technion.ac.il`

MATH DEPARTMENT, TECHNION, HAIFA, 32000, ISRAEL